



Requisition #:

Physician:

Patient Name:

Date of Collection: 3/13/2018

Patient Age: 35

Time of Collection: 07:30 AM

Patient Sex: F

Print Date: 03/21/2018



## Organic Acids Test - Nutritional and Metabolic Profile

Metabolic Markers in Urine      Reference Range (mmol/mol creatinine)      Patient Value      Reference Population - Females Age 13 and Over

### Intestinal Microbial Overgrowth

#### Yeast and Fungal Markers

Marker	Reference Range	Patient Value	Visual Representation
1 Citramalic	≤ 3.6	1.7	Bar chart showing 1.7 within the reference range.
2 5-Hydroxymethyl-2-furoic	≤ 14	13	Bar chart showing 13 within the reference range.
3 3-Oxoglutaric	≤ 0.33	0	Bar chart showing 0.00 within the reference range.
4 Furan-2,5-dicarboxylic	≤ 16	9.5	Bar chart showing 9.5 within the reference range.
5 Furancarboxylglycine	≤ 1.9	1.4	Bar chart showing 1.4 within the reference range.
6 Tartaric	≤ 4.5	3.6	Bar chart showing 3.6 within the reference range.
7 Arabinose	≤ 29	H 194	Bar chart showing 194 significantly above the reference range.
8 Carboxycitric	≤ 29	19	Bar chart showing 19 within the reference range.
9 Tricarballic	≤ 0.44	0.21	Bar chart showing 0.21 within the reference range.

#### Bacterial Markers

10 Hippuric	≤ 613	H 1114	Bar chart showing 1114 significantly above the reference range.
11 2-Hydroxyphenylacetic	0.06 - 0.66	0.31	Bar chart showing 0.31 within the reference range.
12 4-Hydroxybenzoic	≤ 1.3	0.69	Bar chart showing 0.69 within the reference range.
13 4-Hydroxyhippuric	0.79 - 17	10	Bar chart showing 10 within the reference range.
14 DHPA (Beneficial Bacteria)	≤ 0.38	H 2.1	Bar chart showing 2.1 significantly above the reference range.

#### Clostridia Bacterial Markers

15 4-Hydroxyphenylacetic ( <i>C. difficile</i> , <i>C. stricklandii</i> , <i>C. lituseburense</i> & others)	≤ 19	14	Bar chart showing 14 within the reference range.
16 HPPA ( <i>C. sporogenes</i> , <i>C. caloritolerans</i> , <i>C. botulinum</i> & others)	≤ 208	H 350	Bar chart showing 350 significantly above the reference range.
17 4-Cresol ( <i>C. difficile</i> )	≤ 75	20	Bar chart showing 20 within the reference range.
18 3-Indoleacetic ( <i>C. stricklandii</i> , <i>C. lituseburense</i> , <i>C. subterminale</i> & others)	≤ 11	1.6	Bar chart showing 1.6 within the reference range.

Testing performed by The Great Plains Laboratory, Inc., Lenexa, Kansas. The Great Plains Laboratory has developed and determined the performance characteristics of this test. This test has not been evaluated by the U.S. FDA; the FDA does not currently regulate such testing.

# The Great Plains Laboratory, Inc.

Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

Metabolic Markers in Urine      Reference Range (mmol/mol creatinine)      Patient Value      Reference Population - Females Age 13 and Over

## Oxalate Metabolites

19	Glyceric	0.77 - 7.0	3.8	
20	Glycolic	16 - 117	92	
21	Oxalic	6.8 - 101	<b>H</b> 122	

## Glycolytic Cycle Metabolites

22	Lactic	≤ 48	40	
23	Pyruvic	≤ 9.1	5.6	

## Mitochondrial Markers - Krebs Cycle Metabolites

24	Succinic	≤ 9.3	<b>H</b> 10	
25	Fumaric	≤ 0.94	0.86	
26	Malic	0.06 - 1.8	1.3	
27	2-Oxoglutaric	≤ 35	11	
28	Aconitic	6.8 - 28	15	
29	Citric	≤ 507	299	

## Mitochondrial Markers - Amino Acid Metabolites

30	3-Methylglutaric	≤ 0.76	0.31	
31	3-Hydroxyglutaric	≤ 6.2	4.0	
32	3-Methylglutaconic	≤ 4.5	1.3	

## Neurotransmitter Metabolites

### Phenylalanine and Tyrosine Metabolites

33	Homovanillic (HVA) (dopamine)	0.80 - 3.6	3.0	
34	Vanillylmandelic (VMA) (norepinephrine, epinephrine)	0.46 - 3.7	1.0	
35	HVA / VMA Ratio	0.16 - 1.8	<b>H</b> 3.0	

### Tryptophan Metabolites

36	5-Hydroxyindoleacetic (5-HIAA) (serotonin)	≤ 4.3	1.2	
37	Quinolinic	0.85 - 3.9	1.7	
38	Kynurenic	0.17 - 2.2	0.80	
39	Quinolinic / 5-HIAA Ratio	0.42 - 2.0	1.4	

# The Great Plains Laboratory, Inc.

Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

Metabolic Markers in Urine      Reference Range (mmol/mol creatinine)      Patient Value      Reference Population - Females Age 13 and Over

## Pyrimidine Metabolites - Folate Metabolism

40	Uracil	≤ 9.7		3.9	
41	Thymine	≤ 0.56		0.24	

## Ketone and Fatty Acid Oxidation

42	3-Hydroxybutyric	≤ 3.1	<b>H</b>	6.7	
43	Acetoacetic	≤ 10		6.4	
44	4-Hydroxybutyric	≤ 4.8		2.5	
45	Ethylmalonic	0.44 - 2.8		1.7	
46	Methylsuccinic	0.10 - 2.2		1.4	
47	Adipic	0.04 - 3.8		1.2	
48	Suberic	0.18 - 2.2		1.6	
49	Sebacic	≤ 0.24		0.15	

## Nutritional Markers

### Vitamin B12

50	Methylmalonic *	≤ 2.3		1.1	
----	-----------------	-------	--	-----	--

### Vitamin B6

51	Pyridoxic (B6)	≤ 34		8.1	
----	----------------	------	--	-----	--

### Vitamin B5

52	Pantothenic (B5)	≤ 10	<b>H</b>	12	
----	------------------	------	----------	----	--

### Vitamin B2 (Riboflavin)

53	Glutaric *	0.04 - 0.36		0.34	
----	------------	-------------	--	------	--

### Vitamin C

54	Ascorbic	10 - 200		137	
----	----------	----------	--	-----	--

### Vitamin Q10 (CoQ10)

55	3-Hydroxy-3-methylglutaric *	0.17 - 39		13	
----	------------------------------	-----------	--	----	--

### Glutathione Precursor and Chelating Agent

56	N-Acetylcysteine (NAC)	≤ 0.28		0	
----	------------------------	--------	--	---	--

### Biotin (Vitamin H)

57	Methylcitric *	0.19 - 2.7		0.76	
----	----------------	------------	--	------	--

\* A high value for this marker may indicate a deficiency of this vitamin.

# The Great Plains Laboratory, Inc.

Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

Metabolic Markers in Urine      Reference Range (mmol/mol creatinine)      Patient Value      Reference Population - Females Age 13 and Over

## Indicators of Detoxification

### Glutathione

58	Pyroglutamic *	10 - 33	29	
59	2-Hydroxybutyric *	0.03 - 1.8	1.7	

### Ammonia Excess

60	Orotic	0.06 - 0.54	0.31	
----	--------	-------------	------	--

### Aspartame, salicylates, or GI bacteria

61	2-Hydroxyhippuric	≤ 1.3	<b>H</b> 2.1	
----	-------------------	-------	--------------	--

\* A high value for this marker may indicate a Glutathione deficiency.

## Amino Acid Metabolites

62	2-Hydroxyisovaleric	≤ 0.42	0	
63	2-Oxoisovaleric	≤ 2.1	0	
64	3-Methyl-2-oxovaleric	≤ 0.87	0	
65	2-Hydroxyisocaproic	≤ 0.48	0	
66	2-Oxoisocaproic	≤ 0.37	0.23	
67	2-Oxo-4-methylbutyric	≤ 0.16	0.12	
68	Mandelic	≤ 0.21	0.18	
69	Phenyllactic	≤ 0.20	0	
70	Phenylpyruvic	0.20 - 1.9	0.52	
71	Homogentisic	≤ 0.36	0.11	
72	4-Hydroxyphenyllactic	≤ 0.80	0.41	
73	N-Acetylaspartic	≤ 3.0	0.75	
74	Malonic	≤ 9.7	8.9	

## Mineral Metabolism

75	Phosphoric	1 000 - 5 000	3 428	
----	------------	---------------	-------	--

# The Great Plains Laboratory, Inc.

Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

## Indicator of Fluid Intake

76 \*Creatinine 162 mg/dL

\*The creatinine test is performed to adjust metabolic marker results for differences in fluid intake. Urinary creatinine has limited diagnostic value due to variability as a result of recent fluid intake. Samples are rejected if creatinine is below 20 mg/dL unless the client requests results knowing of our rejection criteria.

### Explanation of Report Format

The reference ranges for organic acids were established using samples collected from typical individuals of all ages with no known physiological or psychological disorders. The ranges were determined by calculating the mean and standard deviation (SD) and are defined as  $\pm 2SD$  of the mean. Reference ranges are age and gender specific, consisting of Male Adult ( $\geq 13$  years), Female Adult ( $\geq 13$  years), Male Child ( $< 13$  years), and Female Child ( $< 13$  years).


There are two types of graphical representations of patient values found in the new report format of both the standard Organic Acids Test and the Microbial Organic Acids Test.

The first graph will occur when the value of the patient is within the reference (normal) range, defined as the mean plus or minus two standard deviations.


The second graph will occur when the value of the patient exceeds the upper limit of normal. In such cases, the graphical reference range is "shrunk" so that the degree of abnormality can be appreciated at a glance. In this case, the lower limits of normal are not shown, only the upper limit of normal is shown.

In both cases, the value of the patient is given to the left of the graph and is repeated on the graph inside a diamond. If the value is within the normal range, the diamond will be outlined in black. If the value is high or low, the diamond will be outlined in red.


### Example of Value Within Reference Range



### Example of Elevated Value



## Neurotransmitter Metabolism Markers



The diagram contains the patient's test results for neurotransmitter metabolites and shows their relationship with key biochemical pathways within the axon terminal of nerve cells. The effect of microbial byproducts on the blockage of the conversion of dopamine to norepinephrine is also indicated.

Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

## Interpretation

**High yeast/fungal metabolites (Markers 1,2,3,4,5,6,7,8)** indicate a yeast/fungal overgrowth of the gastrointestinal tract. Prescription or natural (botanical) anti-fungals, along with supplementation of high potency multi-strain probiotics (20-50 billion cfu's), may reduce yeast/fungal levels.

**High hippuric acid (Marker 10)** may derive from food, GI bacterial activity, or exposure to the solvent toluene. Hippuric acid is a conjugate of glycine and benzoic acid formed in the liver. Most hippuric acid in urine is derived from microbial breakdown of chlorogenic acid to benzoic acid. Chlorogenic acid is a common substance in beverages and in many fruits and vegetables, including apples, pears, tea, coffee, sunflower seeds, carrots, blueberries, cherries, potatoes, tomatoes, eggplant, sweet potatoes, and peaches. Benzoic acid is present in high amounts in cranberry juice and is a food preservative. The workplace is the most common source of toluene exposure, but toluene may be absorbed from outgassing of new carpets and other building materials, or absorbed during recreational abuse of solvents such as glue-sniffing. Because most hippuric acid in urine is from GI sources, this marker is a poor indicator of toluene exposure and is being replaced by other markers in occupational safety testing. Bacterial overgrowth can be treated with natural anti-bacterial agents and/or probiotics (30-50 billion cfu's) that include *Lactobacillus rhamnosus*.

**High DHPPA (3,4 dihydroxyphenylpropionic acid) (Marker 14)** indicates excessive intake of chlorogenic acid, a common substance found in beverages and in many fruits and vegetables, including apples, pears, tea, coffee, sunflower seeds, carrots, blueberries, cherries, potatoes, tomatoes, eggplant, sweet potatoes, and peaches. Harmless or beneficial bacteria such as Lactobacilli, Bifidobacteria, and *E. coli* mediate the breakdown of chlorogenic acid to 3,4-dihydroxyphenylpropionic acid (DHPPA), and high values may indicate increased amounts of these species in the GI tract. In addition, one *Clostridia* species, *C. orbiscindens*, can convert the flavanoids luteolin and eriodictyol, occurring only in a relatively small food group that includes parsley, thyme, celery, and sweet red pepper to 3,4-dihydroxyphenylpropionic acid. The quantity of *Clostridia orbiscindens* in the GI tract is negligible (approximately 0.1% of the total bacteria) compared to the predominant flora of *Lactobacilli*, *Bifidobacteria*, and *E. coli*. Consequently, this marker is essentially useless as a general *Clostridia* marker but may be a good indicator of the presence of beneficial flora.

Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

**High HPHPA (3-(3-hydroxyphenyl)-3-hydroxypropionic acid) (Marker 16)** is an abnormal phenylalanine metabolite produced when byproducts of *Clostridium* bacteria combine with human metabolites. High concentrations of this compound cause abnormal behavior by inhibiting metabolism of dopamine to epinephrine, resulting in high levels of the dopamine metabolite homovanillic acid (HVA) in the urine and insufficient epinephrine/norepinephrine in the body. It is associated with behavioral, gastrointestinal, and neuropsychiatric symptoms including tic disorders, depression, autism, schizophrenia, aggression, seizures, anorexia, obsessive compulsive disorder, and hyperactivity. Neuropsychiatric effects are more common when values exceed 500 mmol/mol creatinine.

The *Clostridia* species that cause the greatest quantities of urinary HPHPA are *C. sporogenes*, *C. caloritolerans*, and *C. botulinum*. Additionally, *C. mangenoti*, *C. ghoni*, *C. bifementans*, *C. caproicum*, and *C. sordellii* are also capable of causing elevated urinary levels of HPHPA.

HPHPA precursors are **not** produced by *C. perfringens* -types A-F, *C. tetani*, *C. subterminale*, *C. capitovale*, *C. septicum*, *C. difficile*, *C. histolyticum*, or *C. tertium*.

*C. botulinum* would appear to be an unlikely source unless clinical symptoms of botulism are present. The botulinum toxin can cause a severe [flaccid paralytic](http://en.wikipedia.org/wiki/Flaccid_paralysis) disease in humans and animals and is the most potent toxin known to humankind, with a lethal dose of less than 1 µg in humans. Symptoms of botulism include weakness, impaired vision, fatigue, and impaired speech. This may then be followed by weakness of the arms, chest muscles and legs. Surprisingly, symptoms may sometimes be mild and the severity of symptoms appears to be modulated by the amount of beneficial flora in the intestinal tract. In food borne botulism, symptoms generally begin 18 to 36 hours after eating contaminated food, but they can occur as early as 6 hours or as late as 10 days. *C. caloritolerans* is so named because it can survive at the boiling point for 8 hours. Its extreme resistance to heat may allow common food borne transmission. *C. sporogenes* is the name given to strains of *Clostridium botulinum* that do not produce [botulinum](http://en.wikipedia.org/wiki/Botulinum) neurotoxins. *C. sporogenes* differs from *C. botulinum* by a single gene. *C. sporogenes* is ubiquitous in nature and is commonly found in the flora of humans. *C. sordellii* can be pathogenic and has been implicated in fatal toxic shock syndrome among women of child bearing age.

Treatment with Metronidazole or Vancomycin is almost 100% effective in killing parent *Clostridia* organisms but not their spores. At least three months of probiotic therapy is recommended after antimicrobial treatment due to spore formation by *Clostridia* species. *Clostridia* overgrowth can sometimes be controlled by supplementation with *Lactobacillus rhamnosus* GG (Culturelle) or *Saccharomyces boulardii*. Phenylalanine or tyrosine supplements should be avoided because of the possibility of conversion to HPHPA or other toxic byproducts.

**High oxalic with or without elevated glyceric or glycolic acids (Markers 19,20,21)** may be associated with the genetic hyperoxalurias, autism, women with vulvar pain, fibromyalgia, and may also be due to high vitamin C intake. However, kidney stone formation from oxalic acid was not correlated with vitamin C intake in a very large study. Besides being present in varying concentrations in most vegetables and fruits, oxalates, the mineral conjugate base forms of oxalic acid, are also byproducts of molds such as *Aspergillus* and *Penicillium* and probably *Candida*. If yeast or fungal markers are elevated, antifungal therapy may reduce excess oxalates. High oxalates may cause anemia that is difficult to treat, skin ulcers, muscles pains, and heart abnormalities. Elevated oxalic acid is also the result of anti-freeze (ethylene glycol) poisoning. Oxalic acid is a toxic metabolite of trichloroacetic acid and other environmental pollutants. In addition, decomposing vitamin C may form oxalates during transport or storage.

Elevated oxalate values with a concomitant increase in glycolic acid may indicate genetic hyperoxaluria (type I), whereas increased glyceric acid may indicate a genetic hyperoxaluria (type II). Elevated oxalic acid with normal levels of glyceric or glycolic metabolites rules out a genetic cause for high oxalate. However, elevated oxalates may be due to a new genetic disorder, hyperoxaluria type III.



Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

Regardless of its source, high oxalic acid may contribute to kidney stones and may also reduce ionized calcium. Oxalic acid absorption from the GI tract may be reduced by calcium citrate supplementation before meals. Vitamin B6, arginine, vitamin E, chondroitin sulfate, taurine, selenium, omega-3 fatty acids and/or N-acetyl glucosamine supplements may also reduce oxalates and/or their toxicity. Excessive fats in the diet may cause elevated oxalate if fatty acids are poorly absorbed because of bile salt deficiency. Unabsorbed free fatty acids bind calcium to form insoluble soaps, reducing calcium's ability to bind oxalate and increase its absorption. If taurine is low in a plasma amino acid profile, supplementation with taurine (1000 mg/day) may help stimulate bile salt production (taurocholic acid), leading to better fatty acid absorption and diminished oxalate absorption.

High levels of oxalates are common in autism. Malabsorption of fat and intestinal *Candida* overgrowth are probably the major causes for elevated oxalates in this disorder. Even individuals with elevated glyceric or glycolic acids may not have a genetic disease. To rule out genetic diseases in those people with abnormally high markers characteristic of the genetic diseases, do the following steps: (1) Follow the nutritional steps indicated in this interpretation for one month; (2) If *Candida* is present, treat *Candida* for at least one month; (3) Repeat the organic acid test after abstaining from vitamin C supplements for 48 hours; (4) If the biochemical markers characteristic of genetic oxalate disorders are still elevated in the repeat test, consider DNA tests for the most common mutations of oxalate metabolism. DNA testing for type I hyperoxaluria is available from the Mayo Clinic, Rochester, MN as test #89915 "AGXT Gene, Full Gene Analysis" and, for the p.Gly170Arg mutation only, as # 83643 "Alanine:Glyoxylate Aminotransferase [AGXT] Mutation Analysis [G170R], Blood". Another option to confirm the genetic disease is a plasma oxalate test, also available from the Mayo Clinic (Phone 507.266.5700). Plasma oxalate values greater than 50 micromol/L are consistent with genetic oxalate diseases and may serve as an alternate confirmation test.

Bone tends to be the major repository of excess oxalate in patients with primary hyperoxaluria. Bone oxalate levels are negligible in healthy subjects. Oxalate deposition in the skeleton tends to increase bone resorption and decrease osteoblast activity.

Oxalates may also be deposited in the kidneys, joints, eyes, muscles, blood vessels, brain, and heart and may contribute to muscle pain in fibromyalgia. Oxalate crystal formation in the eyes may be a source of severe eye pain in individuals with autism who may exhibit eye-poking behaviors. High oxalates in the GI tract also may significantly reduce absorption of essential minerals such as calcium, magnesium, zinc, and others.

A low oxalate diet may also be particularly useful in the reduction of body oxalates even if dysbiosis of GI flora is the major source of oxalates. Foods especially high in oxalates include spinach, beets, chocolate, soy, peanuts, wheat bran, tea, cashews, pecans, almonds, berries, and many others. A complete list of high oxalate foods is available online at <http://www.greatplainslaboratory.com/home/eng/oxalates.asp>.

**High succinic acid (Marker 24)** may indicate a relative deficiency of riboflavin and/or coenzyme Q10 (cofactors for succinic dehydrogenase in the Krebs cycle). Supplementation with a minimum of 20 mg riboflavin (which could be provided through a high quality multivitamin) and/or 50 mg/day of coenzyme Q10 is recommended. Clinical observation suggests that succinic acid levels also decrease after treatment for GI dysbiosis.

**VMA levels below the mean (Marker 34)** may indicate lower production of the neurotransmitter norepinephrine or the hormone adrenaline, perhaps due to low dietary intake of the amino acid precursors phenylalanine or tyrosine. Vanilylmandelic acid (VMA) is a metabolite of norepinephrine or adrenaline. Low VMA may also result from blocked conversion of dopamine to norepinephrine by *Clostridia* metabolites. Supplementation with phenylalanine or tyrosine may be beneficial. Enzyme cofactors magnesium, B6 (pyridoxine) or bioperin may also be deficient and respond to supplementation.

**High HVA/VMA ratio (Marker 35)** The most common reason for an elevation of the HVA/VMA ratio is the decreased conversion of dopamine to norepinephrine and epinephrine. The enzyme responsible for this conversion, dopamine beta-hydroxylase, is copper and vitamin C dependent, so an elevated ratio could be due to deficiencies of these cofactors. Another common factor is inhibition of this enzyme by *Clostridia* byproducts. A high HPPHA, 4-Cresol, or other elevations of metabolites would be consistent with the latter explanation.

# The Great Plains Laboratory, Inc.

---

Requisition #:

Physician:

Patient Name:

Date of Collection:

3/13/2018

**5-hydroxyindoleacetic acid (5-HIAA) levels below the mean (Marker 36)** may indicate lower production of the neurotransmitter serotonin. 5-hydroxy-indoleacetic acid is a metabolite of serotonin. Low values have been correlated with symptoms of depression. Supplementation with the precursor 5-HTP (5-hydroxytryptophan) at 50-300 mg/day may be beneficial. Supplementation with tryptophan itself may form the neurotoxic metabolite quinolinic acid, however, 5-HTP is not metabolized to quinolinic acid. Excessive tryptophan supplementation has been associated with eosinophilia myalgia syndrome.

**High 3-hydroxybutyric and/or acetoacetic acids (Markers 42, 43)** indicate increased metabolic utilization of fatty acids. These ketones are associated with diabetes mellitus, fasting, dieting (ketogenic or SCD diet), or illness such as nausea or flu, among many other causes. Regardless of cause, supplementation with L-carnitine or acetyl-L-carnitine (500-1000mg per day) may be beneficial.

**Pyridoxic acid (B6) levels below the mean (Marker 51)** may be associated with less than optimum health conditions (low intake, malabsorption, or dysbiosis). Supplementation with B6 (20 - 50 mg/day) or a multivitamin may be beneficial.

**High pantothenic acid (B5) (Marker 52)** indicates high recent intake of pantothenic acid. Pantothenic acid is an essential B vitamin. Since some individuals may require very high doses of pantothenic acid, high values do not necessarily indicate the need to reduce pantothenic acid intake.

**High 2-hydroxyhippuric acid (Marker 61)** may result after ingestion of aspartame (Nutrasweet®) or salicylates (aspirin), or from GI bacteria converting tyrosine or phenylalanine to salicylic acid. 2-Hydroxyhippuric acid is a conjugate of hydroxybenzoic acid (salicylic acid) and glycine.

**Low values for amino acid metabolites (Markers 62-74)** indicate the absence of genetic disorders of amino acid metabolism. These markers are deamination (ammonia removed) byproducts that are very elevated only when a key enzyme has low activity; slight elevations may indicate a genetic variation or heterozygous condition which may be mitigated with diet or supplementation. Low values are not associated with inadequate protein intake and have not been proven to indicate specific amino acid deficiencies.

High quality nutritional supplements can be purchased through your practitioner or at New Beginnings Nutritionals, [www.NBNUS.com](http://www.NBNUS.com) <<http://www.NBNUS.com>> , or call 877-575-2467.

*The nutritional recommendations in this test are not approved by the US FDA. Supplement recommendations are not intended to treat, cure, or prevent any disease and do not take the place of medical advice or treatment from a healthcare professional.*